



Contents lists available at ScienceDirect

Mutation Research/Genetic Toxicology and Environmental Mutagenesis

journal homepage: www.elsevier.com/locate/gentox
Community address: www.elsevier.com/locate/mutres



Mini review

The role of antioxidants and antioxidant-related enzymes in protective responses to environmentally induced oxidative stress

Jorge Limón-Pacheco, María E. Gensebatt*

Departamento de Medicina Genómica y Toxicología Ambiental, Instituto de Investigaciones Biomédicas,
Universidad Nacional Autónoma de México, Apartado Postal 70-228, 04510, Ciudad Universitaria, México, D.F, Mexico

ARTICLE INFO

Article history:

Received 17 September 2008
Accepted 18 September 2008
Available online 8 October 2008

Keywords:

Environmentally induced oxidative stress
Organ specific response
Antioxidants

ABSTRACT

In aerobic organisms, oxygen is essential for efficient energy production but paradoxically, produces chronic toxic stress in cells. Diverse protective systems must exist to enable adaptation to oxidative environments. Oxidative stress (OS) results when production of reactive oxidative species (ROS) exceeds the capacity of cellular antioxidant defenses to remove these toxic species. Epidemiological and clinical studies have linked environmental factors such as diet and lifestyle to cancer, diabetes, atherosclerosis, and neurodegenerative disorders. All of these conditions, as well as the aging process, are associated with OS due to elevation of ROS or insufficient ROS detoxification. Many environmental pollutants engage signaling pathways that are activated in response to OS. The same sequences of events are also associated with the etiology and early pathology of many chronic diseases. Investigations of oxidative responses in different *in vivo* models suggest that, in complex organisms such as mammals, organs and tissues contain distinct antioxidant systems, and this may form the basis for differential susceptibility to environmental toxic agents. Thus, understanding the pathways leading to the induction of antioxidant responses will enable development of strategies to protect against oxidative damage. We shall review evidence of organ-specific antioxidant responses elicited by environmental pollutants in humans and animal models.

© 2008 Elsevier B.V. All rights reserved.

Contents

1. Introduction	138
1.1. Antioxidants and antioxidant-related enzymes	138
1.2. Organ-specific capacity for antioxidant defense	138
1.3. Metabolism and the formation of reactive species	139
1.4. UV light and ionizing radiation	140
1.5. Metals and metalloids	140
1.6. Pesticides	140
1.7. Halogenated and polycyclic aromatic hydrocarbons (PAHs)	140
1.8. Air particulate matter	141
1.9. Pharmaceutical drugs	141
1.10. Pathological conditions	141
1.11. Basis for sensing environmental oxidative stress and the regulation of antioxidant responses	141
1.12. AP-1, NF- κ B, and Nrf2 as the main transcription factors regulating expression of antioxidants involved in environmental oxidative stress responses	143

Abbreviations: AhR, aryl hydrocarbon receptor; APAP, acetaminophen; DMA, dimethylarsinic acid; ARNT, aryl hydrocarbon nuclear translocator; ASK-1, apoptosis signal-regulating kinase 1; BAP, benzo[a]pyrene; CAT, catalase; CYP, cytochromes P450; EGF, epidermal growth factor; GGT, γ -glutamyltransferase; Grx, glutaredoxin; GSH, glutathione; GSH-Px, glutathione peroxidase; GSSG, oxidized glutathione; GSSG-Rd, glutathione reductase; GST, glutathione transferase; Keap 1, Kelch-like ECH-associated protein 1; MAPK, mitogen activated protein kinases; MDA, malondialdehyde; Nrf2, NF-E2-related factor 2; OS, oxidative stress; PAH, polycyclic aromatic hydrocarbons; PM_{2.5}, particulate matter with a diameter less than 2.5 μ m; PTP, protein tyrosine phosphatase; PTP1B, protein tyrosine phosphatase 1B; ROS, reactive oxidative species; SNO, S-nitrosothiols; SOD, superoxide dismutase; TCDD, tetrachlorodibenzo-p-dioxin; Trx, thioredoxin; TrxR, thioredoxin reductase; γ -GCS, γ -glutamylcysteine synthetase.

* Corresponding author. Tel.: +52 555 622 9179; fax: +52 555 6229182.

E-mail addresses: margen@servidor.unam.mx, mgonsebatt@hotmail.com (M.E. Gensebatt).

2. Concluding remarks.....	144
Conflict of interest.....	144
Acknowledgements.....	144
References.....	144

1. Introduction

In aerobic organisms, oxygen is essential for efficient energy production but paradoxically, produces chronic toxic stress in cells. Thus, protective mechanisms must exist for the removal of toxic oxygen byproducts. Diverse protective systems have evolved to enable adaptation to oxidative environments. These antioxidant defense systems are critical for survival in both prokaryotic and eukaryotic organisms.

Oxidative stress (OS) results when production of reactive oxidative species (ROS) exceeds the capacity of cellular antioxidant defenses to remove these toxic species. Epidemiological and clinical studies have linked environmental factors such as diet and lifestyle (e.g., exposure to ionizing radiation, metals, pesticides, organic toxic persistent compounds, air particles, and certain pharmacological drugs) to cancer, diabetes, atherosclerosis, and neurodegenerative disorders. All of these conditions, as well as the aging process, are associated with OS due to elevation of ROS or insufficient ROS detoxification [1].

Tissues and organs have different rates of metabolic activity and oxygen consumption. Their levels of antioxidants are also different. Such is the case with glutathione (GSH) and cysteine, which are lower in the brain than the liver, kidney, or muscle [2]. Investigations of oxidative responses in different *in vivo* models suggest that, in complex organisms such as mammals, organs and tissues contain distinct antioxidant systems, and this may form the basis for differential susceptibility to environmental toxic agents. Notable advances have been made in our understanding of these distinct systems, with several antioxidant systems and their regulatory pathways being described at the cellular level.

1.1. Antioxidants and antioxidant-related enzymes

Defense mechanisms against free radical-induced oxidative damage include the following: (i) catalytic removal of free radicals and reactive species by factors such as catalase (CAT), superoxide dismutase (SOD), peroxidase, and thiol-specific antioxidants; (ii) binding of proteins (e.g., transferrin, metallothionein, haptoglobins, caeroplasmin) to pro-oxidant metal ions, such as iron and copper; (iii) protection against macromolecular damage by proteins such as stress or heat shock proteins; and (iv) reduction of free radicals by electron donors, such as GSH, vitamin E (α tocopherol), vitamin C (ascorbic acid), bilirubin, and uric acid [2] (Fig. 1).

Animal catalases are heme-containing enzymes that convert hydrogen peroxide (H_2O_2) to water and O_2 , and they are largely localized in subcellular organelles such as peroxisomes. Mitochondria and the endoplasmic reticulum contain little CAT. Thus, intracellular H_2O_2 cannot be eliminated unless it diffuses to the peroxisomes [2]. Glutathione peroxidases (GSH-Px) remove H_2O_2 by coupling its reduction with the oxidation of GSH. GSH-Px can also reduce other peroxides, such as fatty acid hydroperoxides. These enzymes are present in the cytoplasm at millimolar concentrations and also present in the mitochondrial matrix. Most animal tissues contain both CAT and GSH-Px activity.

SODs are metal-containing proteins that catalyze the removal of superoxide, generating water peroxide as a final product of the dismutation. Three isoforms have been identified, and they all are present in all eukaryotic cells. The copper-zinc SOD isoform is

present in the cytoplasm, nucleus, and plasma. On the other hand, the manganese SOD isoform is primarily located in mitochondria.

Dietary micronutrients also contribute to the antioxidant defense system. These include β -carotene, vitamin C, and vitamin E (the vitamin E family comprises both tocopherols and tocotrienols, with α -tocopherol being the predominant and most active form). Water-soluble molecules, such as vitamin C, are potent radical-scavenging agents in the aqueous phase of the cytoplasm, whereas lipid soluble forms, such as vitamin E and β -carotene, act as antioxidants within lipid environments. Selenium, copper, zinc, and manganese are also important elements, since they act as cofactors for antioxidant enzymes. Selenium is considered particularly important in protecting the lipid environment against oxidative injury, as it serves as a cofactor for GSH-Px [2–4].

The most abundant cellular antioxidant is the tripeptide, GSH (L- γ -glutamyl-L-cysteinyl glycine). GSH is synthesized in two steps. First, γ -glutamylcysteine synthetase (γ -GCS) forms a γ -peptide bond between glutamic acid and cysteine, and then GSH synthetase adds glycine. GSH prevents the oxidation of protein thiol groups, either directly by reacting with reactive species or indirectly through glutathione transferases [2–4].

1.2. Organ-specific capacity for antioxidant defense

The GSH system, which includes both thiol (GSH) and disulfide (GSSG) forms of GSH along with related enzymes, provides a good example of the variation in antioxidant capacity that exists among organs. The *de novo* synthesis of GSH mainly occurs in the liver, which contributes almost 90% of GSH under normal physiological conditions [3,4]. In mammals, the concentration of GSH in this organ appears to be the highest among all tissues, with concentrations in hepatocytes ranging from 7 to 10 mM [5]. About 20% of *de novo* synthesized GSH is thought to be exported from rat hepatocytes [6], and it appears to be transported through the canalicular membrane to the bilis, where it reaches concentrations of 8–10 mM [5,7]. In contrast, plasma concentrations of GSH are in the micromolar range [8]. The kidney, spleen, and small intestine have moderate concentrations of GSH, ranging from ~3 to 4 mM [9]. Moderate concentrations (~1–3 mM) are also found in the heart, lung, colon, and brain [9]. In mice, skeletal muscle reportedly contains the highest GSH pool, but this conclusion was based on the total amount of muscular tissue, which makes up almost 40% of body weight in mice [10]. In contrast, other studies have found a concentration of ~1 mM in this tissue [9]. Relative activities of classic antioxidant enzymes, such as γ -glutamyltransferase (GGT), GSH-Px, glutathione reductase (GSSG-Rd), glutathione-S-transferase (GST), and SOD, also appear to be organ-specific. For example, GGT activity is highest in the kidney and lowest in the liver and skeletal muscle [10]. GSH-Px, GST, and SOD activities are higher in liver than the kidney and muscle. GSSG-Rd activity is highest in the kidney, with activity being intermediate in the liver and lowest in muscle [10]. The brain contains only low to moderate activity of SOD, CAT, and GSH-Px as compared to the liver or kidney [11].

Of all the antioxidant systems, the thioredoxin (Trx) system and the GSH-glutaredoxin (GSH-Grx) system are the most important. The Trx system is a thiol-specific antioxidant system that includes NADPH, thioredoxin (Trx), and thioredoxin reductase (TrxR). On




Fig. 1. Air pollution, ionizing radiation, UV light, heavy metals, metalloids, pesticides and polycyclic aromatic hydrocarbons among others, induce the formation of reactive oxygen species (ROS) and reactive nitrogen species (RNS) in target tissues, engaging signaling pathways that are activated in response to oxidative stress (OS). Antioxidant redox systems such as GSH-Grx and Thioredoxin (Trx); antioxidant enzymes like superoxide dismutase (SOD), catalase (CAT) and the action of vitamins E and C neutralize ROS and RNS. OS induce posttranslational modifications in proteins modulating their activities. However, in some cases these posttranslational modifications are also involved in the transcriptional modulation of Phase I and II metabolic enzymes and of antioxidant genes.

the other hand, the GSH-Grx system includes NADPH, GSSG-Rd, GSH/GSSG, and Grx [12]. Interestingly, these antioxidant systems are differentially expressed, at least at the transcriptional level, in adult organs and embryonic stages in mice [13]. In adult mice, the highest level of Trx1 mRNA is found in the kidney, with intermediate levels being present in the lung and the lowest levels being present in the brain, heart, and testis. For Trx2 mRNA, maximal expression occurs in kidney. High levels are also present in the heart and testis, and the lowest levels are present in the lung. In mice, the highest Grx1 mRNA expression occurs in the kidney and the lowest in the testis. Conversely, Grx2 mRNA is highest in the testis, with lower expression in the kidney as well as the brain [13]. Expression patterns differ among not only Grx and Trx redox proteins, but also their corresponding reductases. TrxR1 mRNA is most highly expressed in the liver and kidney. However, TrxR2 mRNA levels are highest in spleen, with slightly lower levels being present in the kidney. Intermediate TrxR2 mRNA levels are present in the liver, heart, and testis, while the lung and brain have the lowest levels. GSSG-Rd mRNA, which is generally more abundant than TrxR1 or TrxR2 mRNA, is most highly expressed in the kidney, and lower levels are present in the lung, liver, testis, and brain. The lowest levels of GSSG-Rd mRNA are found in the spleen and heart [13].

Recent evidence indicates that the organ-specific expression patterns for antioxidant system components may be shared by other factors that are not part of these systems, particularly Bcl-2. Bcl-2 prevents apoptosis through its antioxidant functions [14]. It may do so by regulating cellular GSH levels, γ -GCS expression, and activity of mitogen activated protein kinases (MAPK), Erk1/2, and the transcription factor NF- κ B [15]. Although the *in vivo* organi-

zation of these effects will require further investigation, they might be coordinated through organ-specific expression of Bcl-2. In support of this idea, a study of female CD1 mice revealed that Bcl-2 levels increase from birth to 24 months of age in the lung, liver, heart, kidney, and spleen; however, Bcl-2 is not detected in the brain [16]. According to the authors, Bcl-2 overexpression during aging might be a protective mechanism aimed at preventing cell death. We speculate that this mechanism must be tissue specific, although this hypothesis will require validation in an animal model.

GSH levels as well as the activity and expression of antioxidant enzymes may be organ specific, but they could also be modulated by the metabolic requirements of the tissue [13]. Importantly, each organ contains its own antioxidant capacity. Accordingly, each organ may also have a unique capacity for activation of signaling intermediates.

1.3. Metabolism and the formation of reactive species

Cellular energy metabolism and oxygen consumption are coupled to the generation of ROS. Thus, a reduction in metabolic rate reduces the formation of ROS. This is supported by studies of experimental animals subjected to chronic calorie restriction. There is also limited human evidence from a study performed on a Japanese population in which energy intake was 20% less than the national average. The rates of death due to ROS-associated diseases were decreased in this population, with cerebral vascular disease decreasing by 41%, malignancy by 31%, and heart disease by 41% [17].

The metabolism of toxic compounds could result in the generation of reactive metabolites that have even greater toxicity

and deplete cellular antioxidants. In mammals, up regulation of cytochrome P450 (CYP) has been linked to ROS production [18]. Thus, environmental toxicants are a potential source of ROS. Severe depletion of circulating antioxidants has been observed in smokers [19,20]. In addition, occupational exposure to metals, benzene, cement dust, and multiple other agents is associated with increased lipid peroxidation, increased DNA oxidation, and decreased levels of vitamin E and C [21–25]. CYP monooxygenases are a major source of ROS during ischemia/reperfusion [26]. Accordingly, drugs that inhibit P450 activity protect cells from ROS-induced damage after ischemia or block the formation of catechol estrogens and their subsequent oxidation, leading to decreased oxidative damage [27–29]. Finally, evidence suggests that a higher intake of multiple nutrients, including folic acid, potassium, glucosinolates, diallyl sulfides, and flavonoids, greatly reduces the risk of cardiovascular disease associated with air pollution exposure [30,31].

ROS-induced damage to nucleic acids, proteins, carbohydrates, and lipids alters the function of these macromolecules in cells, tissues, and organs [1,2]. These perturbations elicit adaptive cellular responses that increase antioxidant defenses and repair mechanisms (e.g., DNA repair). Severe oxidative damage to macromolecules leads to cellular death.

Many environmental pollutants engage signaling pathways that are activated in response to OS. The same sequences of events are also associated with the etiology and early pathology of many chronic diseases [32]. Thus, understanding the pathways leading to the induction of antioxidant responses will enable development of strategies to protect against oxidative damage. We shall review evidence of organ-specific antioxidant responses elicited by environmental pollutants in humans and animal models.

1.4. UV light and ionizing radiation

Exposure of skin to UVA (320–400 nm) or UVB (290–320 nm) induces formation of ROS, including the superoxide anion radical ($\bullet\text{O}_2^-$), H_2O_2 , the hydroxyl radical ($\bullet\text{OH}$), singlet oxygen ($^1\text{O}_2$), lipid peroxides (LOOH), lipid peroxide radicals ($\text{LOO}\bullet$), and phase II enzymes modulated by NF-E2-related factor 2 (Nrf2). These reactive species have been linked to skin aging, phototoxicity, inflammation, and malignant tumors [33,34].

One of the main reactive species induced by ionizing radiation is the hydroxyl radical. Ionizing radiation generates the hydroxyl radical through oxidation of water which reacts with cellular components such as sugars, amino acids, phospholipids, DNA bases and organic acids, to produce organic radicals. These secondary ROS may subsequently be converted to hydroxyl radicals through further reduction by cellular metabolic processes such as the Fenton reaction [2].

1.5. Metals and metalloids

Because of its capacity to lose electrons, a metal is primarily thought to be toxic by virtue of its generation of ROS. Thus, exposure to high concentrations of a single heavy metal might result in its accumulation and potentially, oxidative damage [2]. Differential organ responses to metal exposure have been well documented. In several *in vivo* models, heavy metal accumulation correlates with specific organ damage. For example, total parenteral feeding of rats with contaminated solutions produces metalloid (As) and heavy metal deposition (Cr, Pb, Mn, Ge, Sn, and Ba) in the liver, kidney, spleen and brain, but not in the femur, spine, or testis. In addition, pathologic damage occurs only in the liver and kidney of these animals [35], suggesting that important differences in organ susceptibility exist. In rats, high concentrations of Mn and Fe might lead to interaction of these metals during their transfer from the

plasma to the brain, liver, or kidney. The finding that the time course of transfer is slowest in the brain suggests that excessive intake of Fe and Mn may accentuate the risk of tissue damage by Mn in the brain [36]. Similar differences have been found with cadmium chloride in rats [37], with vanadium in mice [38], as well as with As in rabbits and mice [39,40]. In rodents, OS has been observed in multiple organs following exposure to As [41–45], Cd [46–48], Cr [49–51], or Hg [52–54].

1.6. Pesticides

Pesticides are another example of agents that act as pro-oxidants and elicit effects in multiple organs. In some cases, these pro-oxidant effects occur alongside pesticide-induced alterations in target enzymes, many of which participate in neurotransmitter metabolism. For example, paraquat has been extensively studied as an OS inducer, and paraquat toxicity is thought to primarily result from ROS generation and alterations in redox cycling [55]. In rats, paraquat induces alterations in antioxidant systems in many tissues (e.g., liver, blood, kidney, lung), and its targets include GSH, glutathione reductase (GSSG-Rd), CAT, SOD, GSH-Px, and glutathione S-transferases (GST), all of which are traditional biomarkers of OS [56–58]. Malathion, an organophosphorus compound, is another example of a pesticide that induces OS, leading to generation of free radicals and alterations in antioxidant systems in multiple organs in rats [59].

1.7. Halogenated and polycyclic aromatic hydrocarbons (PAHs)

Halogenated aromatic hydrocarbons and PAHs are two of the most extensively studied ubiquitous environmental pollutants, and both cause a broad spectrum of toxic effects in mammals, including carcinogenesis, teratogenesis, and immune dysfunction [60]. Molecular responses to both types of compounds, particularly signal transduction and mechanisms of enzyme activation, have been extensively studied both *in vitro* and *in vivo*. Many of the intracellular signals elicited by these compounds are initiated through the aryl hydrocarbon receptor (AhR)/aryl hydrocarbon nuclear translocator (ARNT) signaling pathway [61]. This signaling pathway includes genes involved in the activation and detoxification of many xenobiotics, such as cytochrome P4501A1, UDP-glucuronosyltransferase, and NADPH quinone oxidoreductase [62,63]. These xenobiotics may generate mutagenic metabolites, DNA strand breaks, ROS, and OS [64,65]. The number of AhR/ARNT-responsive genes known to be involved in cellular processes other than xenobiotic bioactivation has increased [66–68], suggesting that these other processes are influenced by exposure to these compounds.

Of the halogenated aromatic hydrocarbons and PAHs, 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) and benzo[a]pyrene (BAP) are the most ubiquitous and abundant representatives. Increasing evidence in animal models links TCDD and BAP with OS, and these compounds reportedly increase cancer risk in certain organs [69–71]. BAP exposure leads to DNA oxidation, protein oxidation, as well as alterations in SOD and CAT activity in the liver and kidney [69]. BAP administration alone, or together with ethanol, induces changes in GSH levels, malondialdehyde (MDA) levels, and SOD activity in the lung and brain with varying degrees of histological changes [70].

Human exposure to TCDD increases the incidence of cancer, a finding that corroborates its hypothesized association with other health effects, including cardiovascular- and endocrine-related effects [72]. Chen et al. observed increased OS in blood samples from workers exposed to polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans [73]. PAHs are the main carcino-

genic compounds found in urban and indoor air pollution. In Indian children, blood levels of carcinogenic PAHs were significantly associated with increased SOD and CAT activity, but were negatively associated with GSH levels. In these children, high blood levels of total PAHs were also significantly correlated with lipid peroxidation in the blood [74]. Studies also reveal that increased oxidative DNA damage as well as up regulation of genes and proteins involved in OS occurs in individuals exposed to environmental air pollutants such as PAHs, dioxins, and particulate matter [65,75–77]. In a posterior analysis of policeman and bus drivers from three European cities, levels of vitamin A, vitamin E, vitamin C, and folate did not significantly correlate with M1dG adducts (cyclic pyrimidopurine N-1,N² malondialdehyde-2'-deoxyguanosine), 8-oxodG adducts (8-oxo-7,8-dihydro-2'-deoxyguanosine), M1dG adducts with total PAH (bulky), or BAP DNA adducts [78]. Occupational exposure to PAHs has also been associated with oxidative DNA damage [79].

1.8. Air particulate matter

Particulate matter with a diameter less than 2.5 μm (PM_{2.5}) has been linked to increased morbidity and mortality from pulmonary and cardiovascular disease [80]. Animal experiments indicate that ROS are potential mediators of particle-induced cardiovascular effects [81–83]. Studies in humans have shown PM_{2.5} induces the strongest heart rate variability in individuals with inefficient antioxidant responses resulting from combined deletion of glutathione S-transferase M1 (GSTM1) and the presence of ≥ 25 microsatellite (GT)_n repeats in the heme oxygenase-1 (HMOX-1) promoter, which reduce promoter function [84]. In addition, in elderly adults, fine particle exposure elevates exhaled nitric oxide (NO), a biomarker of pulmonary inflammation [85]. Recently, ultrafine PM (<1.8 μm) was shown to induce significantly larger early atherosclerotic lesions in apolipoprotein E-deficient mice than PM_{2.5} or filtered air. In addition, decreased anti-inflammatory capacity of plasma high-density lipoprotein as well as greater systemic OS was observed in these ultrafine PM-exposed animals, as evidenced by a significant increase in hepatic MDA levels and expression of Nrf2-regulated antioxidant genes [86].

1.9. Pharmaceutical drugs

Drug intoxication is often associated with alterations in multiple organs. For example, acetaminophen (APAP), which is widely used as an analgesic and antipyretic, produces severe hepatic-renal damage [87,88] and N-acetyl-p-benzoquinoneimine (NAPQI) metabolite-mediated OS when administered in large doses. The latter is considered the main cause of APAP toxicity. Administration of 250–300 mg kg⁻¹ APAP to mice induces OS in the liver and kidney, as shown by several classic biomarkers including lipid peroxidation products and changes in antioxidant enzyme activities (i.e., SOD, CAT, GST, GSH-Px, cytochrome P450) [87,88]. Conversely, administration of APAP or acetylsalicylic acid to rats inhibits quinolinic acid-induced superoxide generation, lipid peroxidation, and cell damage in the hippocampus [89], suggesting that nervous system responses differ from those of other tissues.

1.10. Pathological conditions

Oxidative stress has been implicated in aging and various pathological conditions including cancer, neurological disorders, diabetes, ischemia/reperfusion [90]. These conditions involve many changes, including alterations in the thiol/disulphide redox state,

impaired glucose tolerance, and activation of inflammatory processes. They are also associated with increased activity of NADPH oxidase or induction of xanthine oxidase with subsequent formation of ROS during chronic inflammation and the generation of damage, particularly during ischemia/reperfusion.

Exposure to agents that increase the formation of ROS might accelerate the initiation of pathological changes. Carcinogenesis models have uncovered a link between an organ's capacity to respond to OS and tumor development. For example, dimethylarsinic acid (DMA), the major metabolite of ingested arsenicals in most mammals, promotes the formation of tumors in the urinary bladder, liver, kidney, and thyroid gland in the rat model [91]. DMA-induced tumor formation in the bladder, skin, and lung might be related to increased DNA oxidation, as evidenced by 8-OHdG formation, and increased cell proliferation observed in these tissues [92,93].

Ambient particulate matter with a diameter <2.5 μm contributes to adverse health effects and is considered a risk factor for the development of ischemic cardiovascular events through the exacerbation of atherosclerosis, coronary artery disease, and the triggering of myocardial infarctions [83]. The smallest particles pose the greater danger because of their high content of organic chemicals and prooxidative potential [86].

Several epidemiological studies have linked chronic exposure to arsenic to type II diabetes [94–101]. The mechanism through which chronic arsenic exposure may produce diabetes is unknown although the contribution of As-induced OS might play a role. It has been documented that decreased uptake of glucose into muscle and adipose tissue under type II diabetes leads to chronic extracellular hyperglycemia, resulting in tissue damage and pathophysiological complications including heart disease, atherosclerosis, cataract formation, peripheral nerve damage, and retinopathy [90,102]. Increased OS has been proposed to be one of the major causes of the hyperglycemia-induced diabetic complications. Under hyperglycemic conditions, ROS formation arises from oxidative phosphorylation and glucose autooxidation, as well as a variety of enzymatic sources such as NADPH oxidase, lipoxygenase, cytochrome P450 monooxygenases, and nitric oxide synthase (NOS) [90].

Neurodegenerative disorders and premature aging due to environmental exposure reflect physiological changes that might be related to free radical formation and multiple organ damage [103,104]. The brain is particularly vulnerable to oxidative damage due to its high oxygen consumption, high content of polyunsaturated fatty acids, and the presence of transition metals as Fe and Cu, which can lead to ROS formation via the Fenton reaction. Oxidative stress increases with age and with exposure to air particulates, arsenic, PAHs, ionizing radiation, etc. [33,34,65,75–77,86,92,104]. Thus exposure can be considered an important causative factor in several neurodegenerative diseases, which are typical in elder individuals [90,103]. The liver, kidneys, heart, and brain of aged rats, compared to those of young rats, contain elevated levels of MDA and reduced levels of GSH, vitamin C, and vitamin E [105]. Lower activities of antioxidant enzymes and increased levels of 8-OHdG are also found in these organs in aged animals [105–108].

1.11. Basis for sensing environmental oxidative stress and the regulation of antioxidant responses

As mentioned earlier, multiple environmental stressors induce OS in tissues, sometimes eliciting distinct responses from each. Depending on the magnitude of the stimulus, these might be adaptive (e.g., modulation of biotransformation mechanisms, induction of antioxidant systems) or they may involve activation of signal

transduction pathways leading to cell death. Thus, production of low levels of ROS or reactive nitrogen species that induce OS without killing cells might serve as a signal for repair, adaptation, survival, or transformation [109].

Cellular responses to OS require a sensor, transducer, and effectors. Identifying these signaling components *in vivo* constitutes a major challenge and raises a big question: How is environmental OS perceived by organisms? Clues are provided by *in vitro* studies, which indicate that cysteine residues play a key physiological role in proteins, allowing them to “sense” redox changes associated with OS.

Studies have demonstrated that ROS induce distinct modifications in cysteine residues, altering protein activity under oxidative conditions. Importantly, this protein “response” depends on the degree of oxidation in the key cysteine residue (e.g., sulfenic, –SOH; sulfinic, –SO₂H and sulfonic, –SO₃H acids). Hence, not all cysteine residues are susceptible to oxidation. Under physiological conditions, the intracellular environment is reduced, and most cysteine residues exist in the form of thiols (SH). Thiols do not normally react with ROS such as H₂O₂, unless the reaction is catalyzed. On the other hand, cysteines in thiolate form (S[–]) do react with H₂O₂, and this reaction could lead to structural and functional changes in the protein (for a complete review see [109]). Examples of proteins that are involved in OS responses and are regulated by cellular redox changes (which are induced by oxidants such as H₂O₂) include thioredoxin [110,111], the transcription factors NF-κB and AP-1 [112–114], peroxiredoxins [115–117], and the negative regulator of Nrf2, Kelch-like ECH-associated protein 1 (Keap1) [118]. Other examples include protein tyrosine phosphatases (PTP) [119], which are also regulated by protein hydroperoxides [120] and the kinase MEKK1 [121], which is negatively modulated by the covalent binding of dietary antioxidants, such as isothiocyanates, to the same regulatory cysteine [122].

H₂O₂ is not the only oxidant molecule involved in redox signaling. Another form of redox regulation, S-nitrosylation, consists of the covalent attachment of NO to protein thiols, resulting in the formation of S-nitrosothiols (SNOs) [123]. Under physiological conditions, SNO protein modification might prevent further cellular oxidative or nitrosative stress [123]. Additionally, this modification is implicated in the regulation of apoptosis, cardiovascular disease, and neurodegenerative disorders [123,124] through regulation of key proteins. For example, in HeLa cells, S-nitrosylation of thioredoxin induces its nuclear translocation, which is related to ERK1/2-mediated survival responses [125]. S-nitrosylation also occurs in caspase-3 and apoptosis signal-regulating kinase 1 (ASK-1), a protein that stimulates apoptosis when activated. S-nitrosylation inhibits both proteins, serving as a negative feedback mechanism that blocks apoptosis [126,127]. S-nitrosothiol-induced inhibition of PTPs such as protein tyrosine phosphatase 1B (PTP1B) could potentially enhance or extend phosphoprotein activity, as in the case of the EGF receptor where activation occurs even in the absence of exogenous EGF [128].

The toxicity of •NO is reflected also in the formation of another covalent posttranslational protein modification named tyrosine nitration (3-NT), which consists in the addition of a nitro group (–NO₂) to a tyrosine residue [129]. This reaction involves peroxynitrite (ONOO[–]), which does not react directly with tyrosine probably due to its short life (10–20 ms), but oxidizes and nitrates through its radical products (i.e. •OH, •NO₂, and CO₃•[–]) [130]. According to some authors, the proteins located proximally to the source of generation of nitrating agents have a higher probability of being nitrated. The secondary structure of the protein, the local environment at the tyrosine residue and the type of nitrating agent influence the selectivity of protein tyrosine nitration [131,132]. In an

in vivo context, protein tyrosine nitration has been observed in several pathological conditions such as neurodegenerative disorders and cardiovascular diseases where inflammatory processes occur [133].

Recently, it has been demonstrated that tyrosine nitration might be involved in signal transduction events as an important posttranslational regulatory modification. For example, ionizing radiation within the therapeutic dose range (5 Gy), causes IκB-α tyrosine 181 nitration leading to dissociation of intact IκB-α from NF-κB as consequence of NOS-1 activation [134]. This suggests that tyrosine nitration might be involved in signal transduction modulation, recently proposed, as another form of posttranslational modification independent of phosphorylation [130,131,134]. In the context of the present work, it would be important to demonstrate a relationship between these posttranslational protein modifications and environmental factors that increase •NO and OS production. This could be the case when exposure is related to inflammatory processes [80].

S-glutathionylation is another mechanism of redox regulation. The formation of mixed disulfides between GSH and cysteinyl residues is a reversible reaction, with reversal being catalyzed by Grx [135]. S-glutathionylation occurs in response not only to moderate oxidative or nitrosative stress, but also under normal conditions, and it appears to be protective under severe OS. S-glutathionylation regulates a wide variety of regulatory, structural, and metabolic proteins that are associated with signaling and metabolic pathways (see [136] for review). Examples of proteins that are negatively regulated by S-glutathionylation include tyrosine hydroxylase [137], PKC-α [138], MEKK1 kinase [121], PTP1B [139,140], and inhibitor of kappa B kinase-β, which inhibits NF-κB [141]. Proteomic analysis has revealed that a variety of proteins with different roles in T lymphocytes are susceptible to S-glutathionylation under OS, suggesting that this posttranslational modification is a global mechanism of redox regulation and may also be of pharmacological importance [142,143].

Little is known about the impact of environmental factors and xenobiotics that induce OS, nitrosative stress, or glutathionylation on redox signaling *in vivo*. Exposure to environmental factors and toxic xenobiotics (e.g., UV light, ionizing radiation, low temperatures, heavy metals) might modulate the redox balance through S-glutathionylation or by activating ROS or NOS signaling [144–147]. In the lung, thioredoxin protects against ROS and modulates the Akt survival pathway following exposure to diesel exhaust particles (DEP) [148]. Akt has a redox-sensitive disulfide, and environmental exposure to DEP might produce an alteration in a key regulatory cysteine. However, more experiments are needed to understand this complex regulation *in vivo*.

Some other aspects, such as the reversibility of this process need more clarification. It is possible that the reversibility of the posttranslational protein modifications depend on the magnitude of the oxidation. In other words, when the amount of modified amino acids is too large, the modification becomes irreversible (Fig. 2).

In this regard, another important question emerges. Is redox signaling similarly modulated in all tissues? Some studies suggest that signaling is tissue-specific. According to Janero et al., redox regulation by nitrosylation must be tissue-specific, since nitrosylation is determined by NO production and metabolic rate [149]. Other studies indicate that differences among species also exist [150]. In mammals, a relationship exists among OS, GSH/GSSG status, formation of nitrosylation products, and life span. This observation introduces the new challenge of identifying specific cellular mechanisms that correlate with *in vivo* changes in the concentration of individual nitrosylation products [151]. We recently demonstrated that, in support of tissue-specific redox signaling, systemic alter-




Fig. 2. Environmental exposure and posttranslational redox modifications of proteins. The posttranslational modifications of proteins probably depend on the magnitude of the oxidant stimulus (dose) and duration of exposure (acute or chronic). A low dose and acute exposure might produce reversible posttranslational modifications (i.e. cysteine oxidation to sulfenic (SOH) or sulfinic (SO₂H) acid), associated to survival, repair or adaptation responses. A high dose or chronic exposure might lead to irreversible posttranslational modification and protein degradation. Events that could cause damage to biomolecules, cell death and altered homeostasis. Similar sequences of events are also associated with the etiology and early pathology of many chronic diseases.

ations in the redox balance maintained by GSH modulate distinct MAPK pathways and gene expression in an organ-specific manner *in vivo* [152]. Environmental toxicant-induced generation of ROS and RNS might modulate the structure and the function of signal transduction proteins and consequently, transcription, allowing cells to respond to environmental stressors through changes in gene expression.

1.12. AP-1, NF- κ B, and Nrf2 as the main transcription factors regulating expression of antioxidants involved in environmental oxidative stress responses

Control of gene expression is a key process that could determine the fate of the cell, leading to survival or death. A considerable amount of evidence demonstrates that NF- κ B and AP-1 play an important role in mediating cellular responses to environmental stressors through their ability to upregulate genes related to numerous cellular processes including cell growth and differentiation, immunity, OS responses, inflammation, and apoptosis. Chemicals, drugs, and other agents that alter the cellular redox status also alter NF- κ B and AP-1 activity [113]. Regulation of NF- κ B appears to be complex, involving more than simple phosphorylation of its subunits (e.g., p65) or dissociation from the I κ B family members [153]. Nevertheless, regulation of NF- κ B is clearly redox-dependent. For example, S-glutathionylation of Cys-62 of the p50 subunit prevents binding of the transcription factor to cognate sites in gene promoters [154]. Also, oxidation of the p50 subunit has recently been shown to be reversed by nuclear and cytoplasmic peroxiredoxin-1 [155]. Reynaert et al. showed that Cys-179 of the beta subunit of the IKK complex undergoes S-glutathionylation, which is reversed by Grx and appears to be responsible for the repression of kinase activity by H₂O₂ [141]. Interestingly, Cys-179 also undergoes S-

nitrosylation and is modulated by redox mechanisms in response to environmental exposure to arsenite [156,157].

AP-1 is a heterodimeric complex of proteins (c-Fos and c-Jun) encoded by the proto-oncogenes *c-fos* and *c-jun*. AP-1 activation is associated with cellular growth, differentiation, neuronal excitation, and stress responses. Indeed, increasing evidence demonstrates a role for AP-1 during OS, as c-Fos and c-Jun are rapidly and transiently induced by a variety of extracellular signals, including agents or factors that alter the redox status. The exact mechanisms responsible for regulation of AP-1 activity are not completely understood. In fact, both oxidants and antioxidants appear to participate in AP-1 regulation. For example, short-term exposure of human fibroblasts to As (III) increases expression of c-Jun and c-Fos, whereas chronic exposure decreases expression of both proteins [158]. Thus, the same toxin can elicit opposing effects depending upon the duration of exposure. Moreover, Abate et al. demonstrated that DNA binding of the Fos-Jun heterodimer is modulated in a redox-sensitive manner through conserved cysteine residues located in the DNA-binding domains of these proteins [159]. Redox-sensitive regulation of c-Jun also occurs through S-glutathionylation of Cys-269 as well as through the formation of an inter-molecular disulfide bridge between cysteine residues near the leucine zipper motif [112]. The AP-1 regulator, redox factor-1 (Ref-1), also undergoes cysteine oxidation, leading to a loss in c-Fos and c-Jun DNA binding activity [159,160].

Both NF- κ B and AP-1 interact *in vivo* in response to environmental toxicant-induced OS. Manna et al. showed that chronic exposure of mice to tobacco smoke increases NF- κ B activity and AP-1 levels in different regions of the brain [161]. Recently, these factors have been shown to have opposite effects on expression of *cyp1a1* mRNA in response to heavy metals and OS, with NF- κ B negatively regulating *cyp1a1* expression [162]. Collectively, these findings suggest

that environmental toxicants might alter redox regulation of these transcription factors, ultimately making differential tissue susceptible to damage.

Cellular responses to environmental and endogenous toxicants involve *de novo* synthesis of many biotransformation and antioxidant enzymes. This synthesis is regulated, in part, by the basic leucine zipper transcription factor, Nrf2. Nrf2 remains inactive in the cytoplasm through its interaction with Keap1, which contains critical cysteine residues within several regions that are susceptible to chemical modification (see [163,164] for reviews). Multiple environmental factors and a variety of chemicals and drugs induce the release of Nrf2 from Keap1, allowing Nrf2 to translocate to the nucleus and activate expression of antioxidant response element (ARE)-containing genes, which include phase I and II genes and genes encoding antioxidants. This response is highly coordinated [152]. Knockout models have demonstrated the importance of Keap1-Nrf2-ARE signaling in distinct tissues. Thus, a disruption in this pathway might affect organ toxicity caused by environmental contaminants [163,164].

2. Concluding remarks

Aerobic organisms express an arsenal of antioxidant enzymes and other protective molecules to maintain a reduced intracellular milieu. Environmentally induced OS modulates the expression and activity of antioxidant enzymes as well as the levels of antioxidant molecules, using pathways similar to those underlying degenerative diseases. Age, gender, genetic susceptibility, diet, and levels of exposure are important factors influencing the capacity of an organism to mount a protective response. Human studies indicate that environmentally induced ROS and RNS elicit antioxidant responses in different tissues. The accumulation of oxidative damage resulting from chronic environmental exposure to PAH, PM, and some metals have been linked to cancer, atherosclerosis, pulmonary disease, and cardiovascular disease. Chemoprevention by nutrients, CYP inhibitors, and antioxidant enzyme inducers has been successfully used to inhibit the formation of reactive species. However, the genetic variables involved in influencing antioxidant responses and metabolic generation of reactive species requires investigation.

To date, evidence suggests that the variation in tissue responses to OS is related to the metabolic capacity of the tissue as well as its antioxidant content. Although a great effort has been made to uncover the role of antioxidant proteins in these responses using knockout animals, new *in vivo* models will still be necessary to unravel the complex interactions among redox signaling mechanisms. Another major challenge will understand how these interactions produce tissue-specific responses within the same organism. Knowledge of specific tissue responses as well as of how these responses are affected by age, gender, genetics, structural complexity of the organ, health condition, and nutritional status will facilitate the development of new clinical strategies for the prevention of cancer and other pathological conditions.

Conflict of interest

None.

Acknowledgements

This work was supported in part by PAPIIT IN 207408 and CONACYT 46341-M. The authors thank Pavel Petrosyan PhD, for his technical help.

References

- [1] H. Sies, Glutathione and its role in cellular functions, *Free Radic. Biol. Med.* 27 (1999) 916–921.
- [2] B. Halliwell, J. Gutteridge (Eds.), *Free Radicals in Biology and Medicine*, Oxford University Press, New York, 1999, pp. 105–245.
- [3] B.H. Lauterburg, J.D. Adams, J.R. Mitchell, Hepatic glutathione homeostasis in the rat: efflux accounts for glutathione turnover, *Hepatology* 4 (1984) 586–590.
- [4] S.M. Deneke, B.L. Fanburg, Regulation of cellular glutathione, *Am. J. Physiol.* 257 (1989) L163–L173.
- [5] C.L. Hammond, R. Marchan, S.M. Krance, N. Ballatori, Glutathione export during apoptosis requires functional multidrug resistance-associated proteins, *J. Biol. Chem.* 282 (2007) 14337–14347.
- [6] H. Sies, A. Wahllander, C.H. Waydhas, in: H. Sies, A. Wendel (Eds.), *Functions of Glutathione in Liver and Kidney*, Springer-Verlag, Berlin, 1978, pp. 120–126.
- [7] N. Ballatori, R. Jacob, J.L. Boyer, Intrabiliary glutathione hydrolysis. A source of glutamate in bile, *J. Biol. Chem.* 261 (1986) 7860–7865.
- [8] A. Pastore, G. Federici, E. Bertini, F. Piemonte, Analysis of glutathione: implication in redox and detoxification, *Clin. Chim. Acta* 333 (2003) 19–39.
- [9] O.W. Griffith, A. Meister, Glutathione: interorgan translocation, turnover, and metabolism, *Proc. Natl. Acad. Sci. U.S.A.* 76 (1979) 5606–5610.
- [10] C. Leeuwenburgh, L.L. Ji, Glutathione depletion in rested and exercised mice: biochemical consequence and adaptation, *Arch. Biochem. Biophys.* 316 (1995) 941–949.
- [11] R. Dringen, Metabolism and functions of glutathione in brain, *Prog. Neurobiol.* 62 (2000) 649–671.
- [12] A. Holmgren, Thioredoxin and glutaredoxin systems, *J. Biol. Chem.* 264 (1989) 13963–13966.
- [13] J. Jurado, M.J. Prieto-Alamo, J. Madrid-Rísquez, C. Pueyo, Absolute gene expression patterns of thioredoxin and glutaredoxin redox systems in mouse, *J. Biol. Chem.* 278 (2003) 45546–45554.
- [14] J.H. Jang, Y.J. Surh, Potentiation of cellular antioxidant capacity by Bcl-2: implications for its antiapoptotic function, *Biochem. Pharmacol.* 66 (2003) 1371–1379.
- [15] J.H. Jang, Y.J. Surh, Bcl-2 attenuation of oxidative cell death is associated with up-regulation of gamma-glutamylcysteine ligase via constitutive NF-kappaB activation, *J. Biol. Chem.* 279 (2004) 38779–38786.
- [16] H. López-Araiza, J.L. Ventura, N.E. López-Diazguerrero, H. González-Marquez, M.C. Gutiérrez-Ruiz, A. Zentella, M. Königsberg, Organ- and tissue-specific alterations in the anti-apoptotic protein Bcl-2 in CD1 female mice of different ages, *Biogerontology* 7 (2006) 63–67.
- [17] L.K. Heilbronn, E. Ravussin, Calorie restriction and aging: review of the literature and implications for studies in humans, *Am. J. Clin. Nutr.* 78 (2003) 361–369.
- [18] M. Paolini, P. Perocco, D. Canistro, L. Valgimigli, G.F. Pedulli, R. Iori, C.D. Croce, G. Cantelli-Forti, M.S. Legator, S.Z. Abdel-Rahman, Induction of cytochrome P450, generation of oxidative stress and *in vitro* cell-transforming and DNA-damaging activities by glucoraphanin, the bioprecursor of the chemopreventive agent sulforaphane found in broccoli, *Carcinogenesis* 25 (2004) 61–67.
- [19] M. Dietrich, G. Block, E.P. Norkus, M. Hudes, M.G. Traber, C.E. Cross, L. Packer, Smoking and exposure to environmental tobacco smoke decrease some plasma antioxidants and increase gamma-tocopherol *in vivo* after adjustment for dietary antioxidant intakes, *Am. J. Clin. Nutr.* 77 (2003) 160–166.
- [20] R.J. Bloomer, Decreased blood antioxidant capacity and increased lipid peroxidation in young cigarette smokers compared to nonsmokers: impact of dietary intake, *Nutr. J.* 6 (2007) 39, <http://www.nutritionj.com/content/6/1/39>.
- [21] J.Y. Kim, S. Mukherjee, L. Ngo, D.C. Christiani, Urinary 8-hydroxy-2'-deoxyguanosine as a biomarker of oxidative DNA damage in workers exposed to fine particulates, *Environ. Health Perspect.* 112 (2004) 666–671.
- [22] S. Aydin, I. Aral, N. Kilic, I. Bakan, S. Aydin, F. Ertan, The level of antioxidant enzymes, plasma vitamins C and E in cement plant workers, *Clin. Chim. Acta* 341 (2004) 193–198.
- [23] P. Rossner Jr., V. Svecovaa, A. Milcovaa, Z. Lnenickovaa, I. Solanskaya, R.J. Sram, Seasonal variability of oxidative stress markers in city bus drivers. Part II. Oxidative damage to lipids and proteins, *Mutat. Res.* 642 (2008) 21–27.
- [24] T. Chia, C.Y. Hsu, H.L. Chen, Oxidative damage of workers in secondary metal recovery plants affected by smoking status and joining the smelting work, *Ind. Health* 46 (2008) 174–182.
- [25] N. Yoshioka, H. Nakashima, K. Hosoda, Y. Eitaki, N. Shimada, K. Omae, Urinary excretion of an oxidative stress marker, 8-hydroxyguanine (8-OH-Gua), among nickel-cadmium battery workers, *J. Occup. Health* 50 (2008) 229–235.
- [26] R.A. Gottlieb, Cytochrome P450: major player in reperfusion injury, *Arch. Biochem. Biophys.* 420 (2003) 262–267.
- [27] E.R. Gross, K. Nithipatikom, A.K. Hsu, J.N. Peart, J.R. Falck, W.B. Campbell, G.J. Gross, Cytochrome P450 omega-hydroxylase inhibition reduces infarct size during reperfusion via the sarcolemmal KATP channel, *J. Mol. Cell. Cardiol.* 37 (2004) 1245–1249.
- [28] Z.H. Chen, Y.J. Hurh, H.K. Na, J.H. Kim, Y.J. Chun, D.H. Kim, K.S. Kang, M.H. Cho, Y.J. Surh, 1,5 resveratrol inhibits TCDD-induced expression of CYP1A1 and CYP1B1 and catechol estrogen-mediated oxidative DNA damage in cultured human mammary epithelial cells, *Carcinogenesis* 25 (2004) 2005–2013.

- [29] M. Khan, I.K. Mohan, V.K. Kutala, D. Kumbala, P. Kuppusamy, Cardioprotection by sulfaphenazole, a cytochrome P450 inhibitor: mitigation of ischemia-reperfusion injury by scavenging of reactive oxygen species, *J. Pharm. Exp. Ther.* 323 (2007) 813–821.
- [30] H.C. Hung, K.J. Josphipura, R. Jiang, F.B. Hu, D. Hunter, S.A. Smith-Warner, G.A. Colditz, B. Rosner, D. Spiegelman, W.C. Willett, Fruit and vegetable intake and risk of major chronic disease, *J. Natl. Cancer Inst.* 96 (2004) 1577–1584.
- [31] A.H. Auchincloss, A.V. Roux, J.T. Dvornch, P.L. Brown, R.G. Barr, M.L. Daviglus, D.C. Goff, J.D. Kaufman, M.S. O'Neill, Associations between recent exposure to ambient fine particulate matter and blood pressure in the multi-ethnic study of atherosclerosis (MESA), *Environ. Health Perspect.* 116 (2008) 486–491.
- [32] B. Hennig, A.S. Ettinger, R.J. Jandacek, S. Koo, C. McClain, H. Seifried, A. Silverstone, B. Watkins, W.A. Suk, Using nutrition for intervention and prevention against environmental chemical toxicity and associated diseases, *Environ. Health Perspect.* 115 (2007) 493–495.
- [33] H. Sakurai, H. Yasui, Y. Yamada, H. Nishimura, M. Shigemoto, Detection of reactive oxygen species in the skin of live mice and rats exposed to UVA light: a research review on chemiluminescence and trials for UVA protection, *Photochem. Photobiol. Sci.* 4 (2005) 715–720.
- [34] L. Marrot, C. Jones, P. Perez, J.R. Meunier, The significance of Nrf2 pathway in (photo)-oxidative stress response in melanocytes and keratinocytes of the human epidermis, *Pigment Cell Melanoma Res.* 21 (2008) 79–88.
- [35] A.L. Buchman, M. Neely, Grossie F.V.B. Jr., L. Truong, E. Lykissa, C. Ahn, Organ heavy-metal accumulation during parenteral nutrition is associated with pathologic abnormalities in rats, *Nutrition* 17 (2001) 600–606.
- [36] A.C. Chua, E.H. Morgan, Effects of iron deficiency and iron overload on manganese uptake and deposition in the brain and other organs of the rat, *Biol. Trace Elem. Res.* 55 (1996) 39–54.
- [37] O. Andersen, J.B. Nielsen, P. Svendsen, Oral cadmium chloride intoxication in mice: effects of dose on tissue damage, intestinal absorption and relative organ distribution, *Toxicology* 48 (1988) 225–236.
- [38] M. Radike, D. Warshawsky, J. Caruso, R. Goth-Goldstein, R. Reilman, T. Collins, M. Yaeger, J. Wang, N. Vela, L. Olsen, J. Schneider, Distribution and accumulation of a mixture of arsenic, cadmium, chromium, nickel, and vanadium in mouse small intestine, kidneys, pancreas, and femur following oral administration in water or feed, *J. Toxicol. Environ. Health A* 65 (2002) 2029–2052.
- [39] C.J. Lin, M.H. Wu, Y.M. Hsueh, S.S. Sun, A.L. Cheng, Tissue distribution of arsenic species in rabbits after single and multiple parenteral administration of arsenic trioxide: tissue accumulation and the reversibility after washout are tissue-selective, *Cancer Chemother. Pharmacol.* 55 (2005) 170–178.
- [40] M.F. Hughes, E.M. Kenyon, B.C. Edwards, C.T. Mitchell, L.M. Razo, D.J. Thomas, Accumulation and metabolism of arsenic in mice after repeated oral administration of arsenate, *Toxicol. Appl. Pharmacol.* 191 (2003) 202–210.
- [41] S. Bhadauria, S.J. Flora, Response of arsenic-induced oxidative stress, DNA damage, and metal imbalance to combined administration of DMSA and monoisoamyl-DMSA during chronic arsenic poisoning in rats, *Cell. Biol. Toxicol.* 23 (2007) 91–104.
- [42] S.J. Flora, S. Bhadauria, S.C. Pant, R.K. Dhaked, Arsenic induced blood and brain oxidative stress and its response to some thiol chelators in rats, *Life Sci.* 77 (2005) 2324–2337.
- [43] R. Gupta, D.K. Dubey, G.M. Kannan, S.J. Flora, Concomitant administration of *Moringa oleifera* seed powder in the remediation of arsenic-induced oxidative stress in mouse, *Cell Biol. Int.* 31 (2007) 44–56.
- [44] M. Mittal, S.J. Flora, Effects of individual and combined exposure to sodium arsenite and sodium fluoride on tissue oxidative stress, arsenic and fluoride levels in male mice, *Chem. Biol. Interact.* 162 (2006) 128–139.
- [45] D. Nandi, R.C. Patra, D. Swarup, Effect of cysteine, methionine, ascorbic acid and thiamine on arsenic-induced oxidative stress and biochemical alterations in rats, *Toxicology* 211 (2005) 26–35.
- [46] S. Nemmiche, D. Chabane-Sari, P. Guiraud, Role of alpha-tocopherol in cadmium-induced oxidative stress in Wistar rat's blood, liver and brain, *Chem. Biol. Interact.* 170 (2007) 221–230.
- [47] N. Yadav, S. Khandelwal, Effect of Picroliv on cadmium-induced hepatic and renal damage in the rat, *Hum. Exp. Toxicol.* 25 (2006) 581–591.
- [48] S. Amara, H. Abdelmelek, C. Garrel, P. Guiraud, T. Douki, J.L. Ravanat, A. Favier, M. Sakly, K. Ben Rhouma, Influence of static magnetic field on cadmium toxicity: study of oxidative stress and DNA damage in rat tissues, *J. Trace Elem. Med. Biol.* 20 (2006) 263–269.
- [49] D. Bagchi, J. Balmoori, M. Bagchi, X. Ye, C.B. Williams, S.J. Stohs, Comparative effects of TCDD, endrin, naphthalene and chromium (VI) on oxidative stress and tissue damage in the liver and brain tissues of mice, *Toxicology* 175 (2002) 73–82.
- [50] X.F. Wang, M.L. Xing, Y. Shen, X. Zhu, L.H. Xu, Oral administration of Cr (VI) induced oxidative stress, DNA damage and apoptotic cell death in mice, *Toxicology* 228 (2006) 16–23.
- [51] A. Scibior, H. Zaporowska, Effects of vanadium (V) and/or chromium (III) on L-ascorbic acid and glutathione as well as iron, zinc, and copper levels in rat liver and kidney, *J. Toxicol. Environ. Health A* 70 (2007) 696–704.
- [52] L. Su, M. Wang, S.T. Yin, H.L. Wang, L. Chen, L.G. Sun, D.Y. Ruan, The interaction of selenium and mercury in the accumulations and oxidative stress of rat tissues, *Ecotoxicol. Environ. Saf.* (2007), PMID: 17644179.
- [53] G. Sener, O. Sehirli, A. Tozan, A. Velioglu-Ovunc, N. Gedik, G.Z. Omurtag, *Ginkgo biloba* extract protects against mercury (II)-induced oxidative tissue damage in rats, *Food Chem. Toxicol.* 45 (2007) 543–550.
- [54] G. Sener, A.O. Sehirli, G. Ayanoglu-Dulger, Melatonin protects against mercury (II)-induced oxidative tissue damage in rats, *Pharmacol. Toxicol.* 93 (2003) 290–296.
- [55] R.J. Dinis-Oliveira, J.A. Duarte, A. Sanchez-Navarro, F. Remião, M.L. Bastos, F. Carvalho, Paraquat poisonings: mechanisms of lung toxicity, clinical features, and treatment, *Crit. Rev. Toxicol.* 8 (2008) 13–71.
- [56] S. Ray, A. Sengupta, A. Ray, Effects of paraquat on anti-oxidant system in rats, *Indian J. Exp. Biol.* 45 (2007) 432–438.
- [57] H. Aoki, Y. Otaka, K. Igarashi, A. Takenaka, Soy protein reduces paraquat-induced oxidative stress in rats, *J. Nutr.* 132 (2002) 2258–2262.
- [58] M. Tomita, H. Katsuyama, T. Okuyama, K. Hidaka, Y. Minatogawa, Changes in gene expression level for defense system enzymes against oxidative stress and glutathione level in rat administered paraquat, *Int. J. Mol. Med.* 15 (2005) 689–693.
- [59] M. Akhgari, M. Abdollahi, A. Kebryaezadeh, R. Hosseini, O. Sabzevari, Biochemical evidence for free radical-induced lipid peroxidation as a mechanism for subchronic toxicity of malathion in blood and liver of rats, *Hum. Exp. Toxicol.* 22 (2003) 205–211.
- [60] A. Kasai, N. Hiramatsu, K. Hayakawa, J. Yao, M. Kitamura, Direct, continuous monitoring of air pollution by transgenic sensor mice responsive to halogenated and polycyclic aromatic hydrocarbons, *Environ. Health Perspect.* 116 (2008) 349–354.
- [61] J.C. Rowlands, J.A. Gustafsson, Aryl hydrocarbon receptor mediated signal transduction, *Crit. Rev. Toxicol.* 27 (1997) 109–134.
- [62] D.W. Nebert, T.P. Dalton, A.B. Okey, F.J. Gonzalez, Role of aryl hydrocarbon receptor-mediated induction of the CYP1 enzymes in environmental toxicity and cancer, *J. Biol. Chem.* 279 (2004) 23847–23850.
- [63] T. Shimada, Xenobiotic-metabolizing enzymes involved in activation and detoxification of carcinogenic polycyclic aromatic hydrocarbons, *Drug Metab. Pharmacokinet.* 21 (2006) 257–276.
- [64] D.W. Nebert, A.L. Roe, M.Z. Dieter, W.A. Solis, Y. Yang, T.P. Dalton, Role of the aromatic hydrocarbon receptor and [Ah] gene battery in the oxidative stress response, cell cycle control, and apoptosis, *Biochem. Pharmacol.* 59 (2000) 65–85.
- [65] R. Singh, B. Kaur, I. Kalina, T.A. Popov, T. Georgieva, S. Garte, B. Binkova, R.J. Sram, E. Taioli, P.B. Farmer, Effects of environmental air pollution on endogenous oxidative DNA damage in humans, *Mutat. Res.* 620 (2007) 71–82.
- [66] Y.C. Cho, W. Zheng, C.R. Jefcoate, Disruption of cell-cell contact maximally but transiently activates AhR-mediated transcription in 10T1/2 fibroblasts, *Toxicol. Appl. Pharmacol.* 199 (2004) 220–238.
- [67] Z. Andrysik, J. Vondracek, M. Machala, P. Krcmar, L. Svihalkova-Sindlerova, A. Kranz, C. Weiss, D. Faust, A. Kozubik, C. Dietrich, The aryl hydrocarbon receptor-dependent deregulation of cell cycle control induced by polycyclic aromatic hydrocarbons in rat liver epithelial cells, *Mutat. Res.* 615 (2007) 87–97.
- [68] F. Castorena-Torres, M. Bermudez de Leon, B. Cisneros, O. Zapata-Perez, J.E. Salinas, A. Albores, Changes in gene expression induced by polycyclic aromatic hydrocarbons in the human cell lines HepG2 and A549, *Toxicol. In Vitro* 22 (2008) 411–421.
- [69] K.B. Kim, B.M. Lee, Oxidative stress to DNA, protein, and antioxidant enzymes (superoxide dismutase and catalase) in rats treated with benzo(a)pyrene, *Cancer Lett.* 113 (1997) 205–212.
- [70] M.H. Emre, A. Aktay, A. Polat, N. Vardt, Effects of benzo(a)pyrene and ethanol on oxidative stress of brain, lung tissues and lung morphology in rats, *Chin. J. Physiol.* 50 (2007) 143–148.
- [71] R. Yoshida, Y. Ogawa, Oxidative stress induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin: an application of oxidative stress markers to cancer risk assessment of dioxins, *Ind. Health* 38 (2000) 5–14.
- [72] P.A. Bertazzi, D. Consonni, S. Bachetti, M. Rubagotti, A. Baccarelli, C. Zocchetti, A.C. Pesatori, Health effects of dioxin exposure: 20-year mortality study, *Am. J. Epidemiol.* 153 (2001) 1031–1044.
- [73] H.L. Chen, C.Y. Hsu, D.Z. Hung, M.L. Hu, Lipid peroxidation and antioxidant status in workers exposed to PCDD/Fs of metal recovery plants, *Sci. Total Environ.* 372 (2006) 12–19.
- [74] V.K. Singh, D.K. Patel, J.S. Ram, N. Mathur, M.K.J. Siddiqui, Blood levels of polycyclic aromatic hydrocarbons in children and their association with oxidative stress indices: an Indian perspective, *Clin. Biochem.* 41 (2008) 152–161.
- [75] S. Loft, H.E. Poulsen, K. Vistisen, L.E. Knudsen, Increased urinary excretion of 8-oxo-2'-deoxyguanosine, a biomarker of oxidative DNA damage in urban bus drivers, *Mutat. Res.* 441 (1999) 11–19.
- [76] B. Marczynski, H.P. Rihs, B. Rossbach, J. Holzer, J. Angerer, M. Scherenberg, G. Hoffmann, T. Bruning, M. Wilhelm, Analysis of 8-oxo-7,8-dihydro-2'-deoxyguanosine and DNA strand breaks in white blood cells of occupationally exposed workers: comparison with ambient monitoring, urinary metabolites and enzyme polymorphisms, *Carcinogenesis* 23 (2002) 273–281.
- [77] M. Kon Kim, S. Oh, J.H. Lee, H. Im, Y.M. Ryu, E. Oh, J. Lee, E. Lee, D. Sul, Evaluation of biological monitoring markers using genomic and proteomic analysis for automobile emission inspectors and waste incinerating workers exposed to polycyclic aromatic hydrocarbons or 2,3,7,8-tetrachlorodibenzo-p-dioxins, *Exp. Mol. Med.* 36 (2004) 396–410.
- [78] R. Singh, R.J. Sram, B. Binkova, I. Kalina, T.A. Popov, T. Georgieva, S. Garte, E. Taioli, P.B. Farmer, The relationship between biomarkers of oxidative DNA damage, polycyclic aromatic hydrocarbon DNA adducts antioxidant status and genetic susceptibility following exposure to environmental air pollution in humans, *Mutat. Res.* 620 (2007) 83–92.

- [79] D. Cavallo, C.L. Ursini, P. Pavazzano, C. Cassinelli, A. Frattini, B. Perniconi, A. Di Francesco, A. Ciervo, B. Rondinone, S. Iavicoli, Sister chromatid exchange and oxidative DNA damage in paving workers exposed to PAHs, *Ann. Hyg.* 50 (2006) 211–218.
- [80] S. Becker, L.A. Dailey, J.M. Soukup, S.C. Grambow, R.B. Devlin, Y.C.T. Huang, Seasonal variations in air pollution particle-induced inflammatory mediator release and oxidative stress, *Environ. Health Perspect.* 113 (2005) 1032–1038.
- [81] R.D. Brook, B. Franklin, W. Cascio, Y. Hong, G. Howard, M. Lipsett, R. Luepker, M. Mittleman, J. Samet, S.C. Smith Jr., I. Tager, Expert panel on population and prevention science of the American Heart Association. Air pollution and cardiovascular disease: a statement for healthcare professionals from the expert panel on population and prevention science of the American Heart Association, *Circulation* 109 (2004) 2655–2671.
- [82] S.A. Gurgueira, J. Lawrence, B. Coull, G.G. Murthy, B. Gonzalez-Flecha, Rapid increases in the steady-state concentration of reactive oxygen species in the lungs and heart after particulate air pollution inhalation, *Environ. Health Perspect.* 110 (2002) 749–755.
- [83] A. Nel, Atmosphere. Air pollution-related illness: effects of particles, *Science* 308 (2005) 804–806.
- [84] T. Chahine, A. Baccarelli, A. Litonjua, R.O. Wright, H. Suh, D.R. Gold, D. Sparrow, P. Vokonas, J. Schwartz, Particulate air pollution, oxidative stress genes, and heart rate variability in an elderly cohort, *Environ. Health Perspect.* 115 (2007) 1617–1622.
- [85] S.D. Adar, G. Adamkiewicz, D.R. Gold, J. Schwartz, B.A. Coull, H. Suh, Ambient and microenvironmental particles and exhaled nitric oxide before and after a group bus trip, *Environ. Health Perspect.* 115 (2007) 507–512.
- [86] J.A. Araujo, B. Barajas, M. Kleinman, X. Wang, B.J. Bennett, K.W. Gong, M. Navab, J. Harkema, C. Sioutas, A.J. Lusis, Ambient particulate pollutants in the ultrafine range promote early atherosclerosis and systemic oxidative stress, *Circ. Res.* 102 (2008) 589–596.
- [87] A. Ghosh, P.C. Sil, Anti-oxidative effect of a protein from *Cajanus indicus* L against acetaminophen-induced hepato-nephro toxicity, *J. Biochem. Mol. Biol.* 40 (2007) 1039–1049.
- [88] M.T. Olaleye, B.T. Rocha, Acetaminophen-induced liver damage in mice: effects of some medicinal plants on the oxidative defense system, *Exp. Toxicol. Pathol.* 59 (2008) 319–327.
- [89] H. Maharaj, D.S. Maharaj, S. Daya, Acetylsalicylic acid and acetaminophen protect against oxidative neurotoxicity, *Metab. Brain Dis.* 21 (2006) 189–199.
- [90] M. Valko, D. Leibfritz, J. Moncol, M.T. Cronin, M. Mazur, J. Telser, Free radicals and antioxidants in normal physiological functions and human disease, *Int. J. Biochem. Cell. Biol.* 39 (2007) 44–84.
- [91] S. Yamamoto, H. Waniuchi, T. Hori, Y. Yano, I. Matsui-Yuasa, S. Otani, H. Chen, K. Yoshida, K. Kuroda, G. Endo, S. Fukushima, Possible carcinogenic potential of dimethylarsinic acid as assessed in rat in vivo models: a review, *Mutat. Res.* 386 (1997) 353–361.
- [92] A. Kinoshita, H. Waniuchi, M. Wei, T. Yunoki, S. Fukushima, Elevation of 8-hydroxydeoxyguanosine and cell proliferation via generation of oxidative stress by organic arsenicals contributes to their carcinogenicity in the rat liver and bladder, *Toxicol. Appl. Pharmacol.* 221 (2007) 295–305.
- [93] M. Mizoi, F. Takabayashi, M. Nakano, Y. An, Y. Sagesaka, K. Kato, S. Okada, K. Yamanaka, The role of trivalent dimethylated arsenic in dimethylarsinic acid-promoted skin and lung tumorigenesis in mice: tumor-promoting action through the induction of oxidative stress, *Toxicol. Lett.* 158 (2005) 87–94.
- [94] M. Rahman, M. Tondel, S.A. Ahmad, O. Axelsson, Diabetes mellitus associated with arsenic exposure in Bangladesh, *Am. J. Epidemiol.* 148 (1998) 198–203.
- [95] C.-H. Tseng, T.-Y. Tai, C.-K. Chong, C.-P. Seng, M.-S. Lai, B.J. Lin, H.-Y. Chiou, Y.-M. Hsueh, K.-H. Hsu, C.-J. Chen, Long-term arsenic exposure and incidence of non-insulin dependent diabetes mellitus: a cohort study in arseniasis hyperendemic villages in Taiwan, *Environ. Health Perspect.* 108 (2000) 847–851.
- [96] S.-L. Wang, J.-M. Chiou, C.-J. Chen, C.-H. Tseng, W.-L. Chou, C.-C. Wang, T.-N. Wu, L.W. Chang, Prevalence of non-insulin-dependent diabetes mellitus and related vascular diseases in southwestern arseniasis-endemic and nonendemic areas in Taiwan, *Environ. Health Perspect.* 111 (2003) 155–159.
- [97] J.A. Coronado-González, L.M. Del Razo, G. Garcia-Vargas, F. Sanmiguel-Salazar, J. Escobedo-de la Peña, Inorganic arsenic exposure and type 2 diabetes mellitus in Mexico, *Environ. Res.* 104 (2007) 383–389.
- [98] A. Navas-Acien, E.K. Silbergeld, R.A. Streeter, J.M. Clark, T.A. Burke, E. Guallar, Arsenic exposure and type 2 diabetes: a systematic review of the experimental and epidemiological evidence, *Environ. Health Perspect.* 114 (2006) 641–648.
- [99] C.-J. Chen, S.-L. Wang, J.-M. Chiou, C.-H. Tseng, H.-Y. Chiou, Y.-M. Hsueh, S.-Y. Chen, M.-M. Wu, M.-S. Lai, Arsenic and diabetes and hypertension in human populations: a review, *Toxicol. Appl. Pharmacol.* 222 (2007) 298–304.
- [100] M.-S. Lai, Y.-M. Hsueh, C.-J. Chen, M.-P. Shyu, S.-Y. Chen, T.-L. Kuo, M.-M. Wu, T.-Y. Tai, Ingested inorganic arsenic and prevalence of diabetes mellitus, *Am. J. Epidemiol.* 139 (1994) 484–492.
- [101] A. Navas-Acien, E.K. Silbergeld, R. Pastor-Barriuso, E. Guallar, Arsenic exposure and prevalence of Type 2 Diabetes in US adults, *JAMA* 300 (2008) 845–846.
- [102] M. Brownlee, A. Cerami, The biochemistry of the complications of diabetes-mellitus, *Ann. Rev. Biochem.* 50 (1981) 385–432.
- [103] V.B. Junqueira, S.B. Barros, S.S. Chan, L. Rodrigues, L. Giavarotti, R.L. Abud, G.P. Deucher, Aging and oxidative stress, *Mol. Aspects Med.* 25 (2004) 5–16.
- [104] G. I Jenkins, Molecular mechanisms of skin ageing, *Mech. Ageing Dev.* 123 (2002) 801–810.
- [105] T. Jayakumar, P.A. Thomas, P. Geraldine, Protective effect of an extract of the oyster mushroom, *Pleurotus ostreatus*, on antioxidants of major organs of aged rats, *Exp. Gerontol.* 42 (2007) 183–191.
- [106] L. Tian, Q. Cai, H. Wei, Alterations of antioxidant enzymes and oxidative damage to macromolecules in different organs of rats during aging, *Free Radic. Biol. Med.* 24 (1998) 1477–1484.
- [107] Y.T. Wong, R. Ruan, F.E. Tay, Relationship between levels of oxidative DNA damage, lipid peroxidation and mitochondrial membrane potential in young and old F344 rats, *Free Radic. Res.* 40 (2006) 393–402.
- [108] M.L. Hamilton, H. Van Remmen, J.A. Drake, H. Yang, Z.M. Guo, K. Kewitt, C.A. Walter, A. Richardson, Does oxidative damage to DNA increase with age? *Proc. Natl. Acad. Sci. U.S.A.* 98 (2001) 10469–10474.
- [109] H.J. Forman, J.M. Fukuto, M. Torres, Redox signaling: thiol chemistry defines which reactive oxygen and nitrogen species can act as second messengers, *Am. J. Physiol. Cell. Physiol.* 287 (2004) C246–C256.
- [110] A. Holmgren, Thioredoxin structure and mechanism: conformational changes on oxidation of the active-site sulfhydryls to a disulfide, *Structure* 3 (1995) 239–243.
- [111] V.V. Sumbayev, S-nitrosylation of thioredoxin mediates activation of apoptosis signal-regulating kinase 1, *Arch. Biochem. Biophys.* 415 (2003) 133–136.
- [112] P. Klatt, E.P. Molina, M.G. De Lacoba, C.A. Padilla, E. Martinez-Galesteo, J.A. Barcena, S. Lamas, Redox regulation of c-Jun DNA binding by reversible S-glutathiolation, *FASEB J.* 13 (1999) 1481–1490.
- [113] D. Gius, A. Botero, S. Shah, H.A. Curry, Intracellular oxidation/reduction status in the regulation of transcription factors NF-kappaB and AP-1, *Toxicol. Lett.* 106 (1999) 93–106.
- [114] T. Nishi, N. Shimizu, M. Hiramoto, I. Sato, Y. Yamaguchi, M. Hasegawa, S. Aizawa, H. Tanaka, K. Kataoka, H. Watanabe, H. Handa, Spatial redox regulation of a critical cysteine residue of NF-kappa B in vivo, *J. Biol. Chem.* 277 (2002) 44548–44556.
- [115] S.W. Kang, T.S. Chang, T.H. Lee, E.S. Kim, D.Y. Yu, S.G. Rhee, Cytosolic peroxiredoxin attenuates the activation of Jnk and p38 but potentiates that of Erk in HeLa cells stimulated with tumor necrosis factor-alpha, *J. Biol. Chem.* 279 (2004) 2535–2543.
- [116] S.G. Rhee, H.Z. Chae, K. Kim, Peroxiredoxins: a historical overview and speculative preview of novel mechanisms and emerging concepts in cell signaling, *Free Radic. Biol. Med.* 38 (2005) 1543–1552.
- [117] W. Jeong, S.J. Park, T.S. Chang, D.Y. Lee, S.G. Rhee, Molecular mechanism of the reduction of cysteine sulfenic acid of peroxiredoxin to cysteine by mammalian sulfiredoxin, *J. Biol. Chem.* 281 (2006) 14400–14407.
- [118] N. Wakabayashi, A.T. Dinkova-Kostova, W.D. Holtzclaw, M.I. Kang, A. Kobayashi, M. Yamamoto, T.W. Kensler, P. Talalay, Protection against electrophile and oxidant stress by induction of the phase 2 response: fate of cysteines of the Keap1 sensor modified by inducers, *Proc. Natl. Acad. Sci. U.S.A.* 101 (2004) 2040–2045.
- [119] J.M. Denu, K.G. Tanner, Specific and reversible inactivation of protein tyrosine phosphatases by hydrogen peroxide: evidence for a sulfenic acid intermediate and implications for redox regulation, *Biochemistry* 37 (1998) 5633–5642.
- [120] M. Gracianin, M.J. Davies, Inhibition of protein tyrosine phosphatases by amino acid, peptide, and protein hydroperoxides: potential modulation of cell signaling by protein oxidation products, *Free Radic. Biol. Med.* 42 (2007) 1543–1551.
- [121] J.V. Cross, D.J. Templeton, Oxidative stress inhibits MEK1 by site-specific glutathionylation in the ATP-binding domain, *Biochem. J.* 381 (2004) 675–683.
- [122] J.V. Cross, F.W. Foss, J.M. Rady, T.L. Macdonald, D.J. Templeton, The isothiocyanate class of bioactive nutrients covalently inhibits the MEK1 protein kinase, *BMC Cancer* 7 (2007) 183.
- [123] J. Sun, C. Steenbergen, E. Murphy, S-nitrosylation: NO-related redox signaling to protect against oxidative stress, *Antioxid. Redox Signal.* 8 (2006) 1693–1705.
- [124] K.K. Chung, T.M. Dawson, V.L. Dawson, Nitric oxide, S-nitrosylation and neurodegeneration, *Cell Mol. Biol.* 51 (2005) 247–254.
- [125] R.J. Arai, H. Masutani, J. Yodoi, V. Debbas, F.R. Laurindo, A. Stern, H.P. Monteiro, Nitric oxide induces thioredoxin-1 nuclear translocation: possible association with the p21Ras survival pathway, *Biochem. Biophys. Res. Commun.* 348 (2006) 1254–1260.
- [126] D.A. Mitchell, M.A. Marletta, Thioredoxin catalyzes the S-nitrosation of the caspase-3 active site cysteine, *Nat. Chem. Biol.* 1 (2005) 154–158.
- [127] J.B. Mannick, Regulation of apoptosis by protein S-nitrosylation, *Amino Acids* 32 (2007) 523–526.
- [128] S. Li, A.R. Whorton, Regulation of protein tyrosine phosphatase 1B in intact cells by S-nitrosothiols, *Arch. Biochem. Biophys.* 410 (2003) 269–279.
- [129] M. Souza, G. Peluffo, R. Radi, Protein tyrosine nitration-functional alteration or just a biomarker? *Free Radic. Biol. Med.* 45 (2008) 357–366.
- [130] R. Radi, Nitric oxide, oxidants, and protein tyrosine nitration, *Proc. Natl. Acad. Sci. U.S.A.* 101 (2004) 4003–4008.
- [131] H. Ischiropoulos, Biological selectivity and functional aspects of protein tyrosine nitration, *Biochem. Biophys. Res. Commun.* 305 (2003) 776–783.
- [132] J.M. Souza, E. Daikhin, M. Yudkoff, C.S. Raman, H. Ischiropoulos, Factors determining the selectivity of protein tyrosine nitration, *Arch. Biochem. Biophys.* 371 (1999) 169–178.
- [133] H. Ischiropoulos, Biological tyrosine nitration: a pathophysiological function of nitric oxide and reactive oxygen species, *Arch. Biochem. Biophys.* 356 (1998) 1–11.
- [134] V.A. Yakovlev, I.J. Barani, C.S. Rabender, S.M. Black, J.K. Leach, P.R. Graves, G.E. Kellogg, R.B. Mikkelsen, Tyrosine nitration of IkappaBalpha: a novel

- mechanism for NF-kappaB activation, *Biochemistry* 46 (2007) 11671–11683.
- [135] M.M. Gallogly, J.J. Mielay, Mechanisms of reversible protein glutathionylation in redox signaling and oxidative stress, *Curr. Opin. Pharmacol.* 7 (2007) 381–391.
- [136] I. Dalle-Donne, R. Rossi, D. Giustarini, R. Colombo, A. Milzani, S-glutathionylation in protein redox regulation, *Free Radic. Biol. Med.* 43 (2007) 883–898.
- [137] C.R. Borges, T. Geddes, J.T. Watson, D.M. Kuhn, Dopamine biosynthesis is regulated by S-glutathionylation: potential mechanism of tyrosine hydroxylase inhibition during oxidative stress, *J. Biol. Chem.* 277 (2002) 48295–48302.
- [138] N.E. Ward, J.R. Stewart, C.G. Ioannides, C.A. O'Brian, Oxidant-induced S-glutathiolation inactivates protein kinase C-alpha (PKC-alpha): a potential mechanism of PKC isozyme regulation, *Biochemistry* 39 (2000) 10319–10329.
- [139] S.R. Lee, K.S. Kwon, S.R. Kim, S.G. Rhee, Reversible inactivation of protein-tyrosine phosphatase 1B in A431 cells stimulated with epidermal growth factor, *J. Biol. Chem.* 273 (1998) 15366–15372.
- [140] W.C. Barrett, J.P. DeGnore, S. König, H.M. Fales, Y.F. Keng, Z.Y. Zhang, M.B. Yim, P.B. Chock, Regulation of PTP1B via glutathionylation of the active site cysteine 215, *Biochemistry* 38 (1999) 6699–6705.
- [141] N.L. Reynaert, A. van der Vliet, A.S. Guala, T. McGovern, M. Hristova, C. Pantano, N.H. Heintz, J. Heim, Y.S. Ho, D.E. Matthews, E.F. Wouters, Y.M. Janssen-Heininger, Dynamic redox control of NF-kappaB through glutaredoxin-regulated S-glutathionylation of inhibitory kappaB kinase beta, *Proc. Natl. Acad. Sci. U.S.A.* 103 (2006) 13086–13091.
- [142] M. Fratelli, H. Demol, M. Puype, S. Casagrande, I. Eberini, M. Salmona, V. Bonetto, M. Mengozzi, F. Duffieux, E. Miclet, A. Bachi, J. Vandekerckhove, E. Gianazza, P. Ghezzi, Identification by redox proteomics of glutathionylated proteins in oxidatively stressed human T lymphocytes, *Proc. Natl. Acad. Sci. U.S.A.* 99 (2002) 3505–3510.
- [143] P. Ghezzi, P. Di Simplicio, Glutathionylation pathways in drug response, *Curr. Opin. Pharmacol.* 7 (2007) 398–403.
- [144] G.S. Buzard, K.S. Kasprzak, Possible roles of nitric oxide and redox cell signaling in metal-induced toxicity and carcinogenesis: a review, *J. Environ. Pathol. Toxicol. Oncol.* 19 (2000) 179–199.
- [145] J.B. Mannick, C.M. Schonhoff, Analysis of protein S-nitrosylation, *Curr. Protoc. Protein Sci.* (2006), Unit 14.6 (Chapter 14).
- [146] K. Kato, M. Mizoi, Y. An, M. Nakano, H. Wanibuchi, G. Endo, Y. Endo, M. Hoshino, S. Okada, K. Yamanaka, Oral administration of diphenylarsinic acid, a degradation product of chemical warfare agents, induces oxidative and nitrosative stress in cerebellar Purkinje cells, *Life Sci.* 81 (2007) 1518–1525.
- [147] D.P. Blagojević, Antioxidant systems in supporting environmental and programmed adaptations to low temperatures, *Cryo Lett.* 28 (2007) 137–150.
- [148] M. Kaimul Ahsan, H. Nakamura, M. Tanito, K. Yamada, H. Utsumi, J. Yodoi, Thioredoxin-1 suppresses lung injury and apoptosis induced by diesel exhaust particles (DEP) by scavenging reactive oxygen species and by inhibiting DEP-induced downregulation of Akt, *Free Radic. Biol. Med.* 39 (2005) 1549–1559.
- [149] D.R. Janero, N.S. Bryan, F. Saijo, V. Dhawan, D.J. Schwalb, M.C. Warren, M. Feelisch, Differential nitrosylation of blood and tissue constituents during glycyl trinitrate biotransformation in vivo, *Proc. Natl. Acad. Sci. U.S.A.* 101 (2004) 16958–16963.
- [150] M. Feelisch, T. Rassaf, S. Mnaimneh, N. Singh, N.S. Bryan, D. Jourdain, M. Kelm, Concomitant S-, N-, and heme-nitrosylation in biological tissues and fluids: implications for the fate of NO in vivo, *FASEB J.* 16 (2002) 1775–1785.
- [151] N.S. Bryan, T. Rassaf, R.E. Maloney, C.M. Rodriguez, F. Saijo, J.R. Rodriguez, M. Feelisch, Cellular targets and mechanisms of nitrosylation: an insight into their nature and kinetics in vivo, *Proc. Natl. Acad. Sci. U.S.A.* 101 (2004) 4308–4313.
- [152] J.H. Limón-Pacheco, N.A. Hernández, M.L. Fanjul-Moles, M.E. Gonsébat, Glutathione depletion activates mitogen-activated protein kinase (MAPK) pathways that display organ-specific responses and brain protection in mice, *Free Radic. Biol. Med.* 43 (2007) 1335–1347.
- [153] Y.M. Janssen-Heininger, M.E. Poynter, P.A. Baeuerle, Recent advances towards understanding redox mechanisms in the activation of nuclear factor kappaB, *Free Radic. Biol. Med.* 28 (2000) 1317–1327.
- [154] E. Pineda-Molina, P. Klatt, J. Vázquez, A. Marina, M. García de Lacoba, D. Pérez-Sala, S. Lamas, Glutathionylation of the p50 subunit of NF-kappaB: a mechanism for redox-induced inhibition of DNA binding, *Biochemistry* 40 (2001) 14134–14142.
- [155] J.M. Hansen, S. Moriarty-Craige, D.P. Jones, Nuclear and cytoplasmic peroxiredoxin-1 differentially regulates NF-kappaB activities, *Free Radic. Biol. Med.* 43 (2007) 282–288.
- [156] P. Kapahi, T. Takahashi, G. Natoli, S.R. Adams, Y. Chen, R.Y. Tsien, M. Karin, Inhibition of NF-kappa B activation by arsenite through reaction with a critical cysteine in the activation loop of I kappa B kinase, *J. Biol. Chem.* 275 (2000) 36062–36066.
- [157] N.L. Reynaert, K. Kless, S.H. Korn, N. Vos, A.S. Guala, E.F. Wouters, A. van der Vliet, Y.M. Janssen-Heininger, Nitric oxide represses inhibitory kappaB kinase through S-nitrosylation, *Proc. Natl. Acad. Sci. U.S.A.* 101 (2004) 8945–8950.
- [158] Y. Hu, X. Jin, E.T. Snow, Effect of arsenic on transcription factor AP-1 and NF-kappaB DNA binding activity and related gene expression, *Toxicol. Lett.* 133 (2002) 33–45.
- [159] C. Abate, L. Patel, F.J. Rauscher, T. Curran, Redox regulation of fos and jun DNA-binding activity in vitro, *Science* 249 (1990) 1157–1161.
- [160] S. Xanthoudakis, G. Miao, F. Wang, Y.C. Pan, T. Curran, Redox activation of Fos-Jun DNA binding activity is mediated by a DNA repair enzyme, *EMBO J.* 11 (1992) 3323–3335.
- [161] S.K. Manna, T. Rangasamy, K. Wise, S. Sarkar, S. Shishodia, S. Biswal, G.T. Ramesh, Long term environmental tobacco smoke activates nuclear transcription factor-kappa B, activator protein-1, and stress responsive kinases in mouse brain, *Biochem. Pharmacol.* 71 (2006) 1602–1609.
- [162] H.M. Korashy, A.O. El-Kadi, The role of redox-sensitive transcription factors NF-kappaB and AP-1 in the modulation of the Cyp1a1 gene by mercury, lead, and copper, *Free Radic. Biol. Med.* 44 (2008) 795–806.
- [163] T.W. Kensler, N. Wakabayashi, S. Biswal, Cell survival responses to environmental stresses via the Keap1-Nrf2-ARE pathway, *Annu. Rev. Pharmacol. Toxicol.* 47 (2007) 89–116.
- [164] I.M. Copple, C.E. Goldring, N.R. Kitteringham, B.K. Park, The Nrf2-Keap1 defence pathway: role in protection against drug-induced toxicity, *Toxicology* 246 (2008) 24–33.